

The anisotropic flow of thermal photons in most-central α -clustered C+Au collisions at 200A GeV

Tuesday, 15 November 2022 15:10 (20)

The light nuclei such as B, C, O etc. can occur in different stable α -clustered states, as found in the low-energy nuclear structure studies. Recent phenomenological findings suggest that such light nuclei with different exotic shapes can produce large initial-state anisotropies in relativistic nuclear collisions. The electromagnetic radiations, being sensitive to the initial dynamics, are expected to be affected by the initial clustered structures significantly. In this work, we estimate the production and anisotropic flow of photons from most-central collisions of triangular α -clustered carbon and gold at $\sqrt{s_{NN}} = 200$ GeV at RHIC using an event-by-event hydrodynamic framework and compare the results with those obtained from unclustered carbon and gold collisions. We find that the thermal photon v_3 for most central collisions is significantly large for the clustered case compared to the case with unclustered carbon. In contrast, the elliptic flow parameter (v_2) is found to be similar for the two cases. We show that the ratio of anisotropic flow coefficients can be a potential observable to detect the α -clustered structure in the carbon nucleus.

Primary author(s) : Dr DASGUPTA, Pingal (Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University); Dr CHATTERJEE, Rupa (Variable Energy Cyclotron Centre, 1/AF, Bidhan Nagar, Kolkata-700064, India); Dr MA, Guo-Liang (Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University, Shanghai 200433, China)

Presenter(s) : Dr DASGUPTA, Pingal (Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University)

Session Classification : Day 1: Session 3

Track Classification : Electromagnetic probes